

# THE $SL(2)$ -TYPE AND BASE CHANGE

OMER OFFEN AND EITAN SAYAG

ABSTRACT. The  $SL(2)$ -type of any smooth, irreducible and unitarizable representation of  $GL_n$  over a  $p$ -adic field was defined by Venkatesh. We provide a natural way to extend the definition to all smooth and irreducible representations. For unitarizable representations we show that the  $SL(2)$ -type of a representation is preserved under base change with respect to any finite extension. The Klyachko model of a smooth, irreducible and unitarizable representation  $\pi$  of  $GL_n$  depends only on the  $SL(2)$ -type of  $\pi$ . As a consequence we observe that the Klyachko model of  $\pi$  and of its base-change are of the same type.

## 1. INTRODUCTION

Let  $F$  be a finite extension of  $\mathbb{Q}_p$ . In [Ven05], Venkatesh assigned a partition of  $n$ , the  $SL(2)$ -type of  $\pi$ , to any smooth, irreducible and unitarizable representation  $\pi$  of  $GL_n(F)$ . For a representation of Arthur type the  $SL(2)$ -type encodes the combinatorial data in the Arthur parameter. In general, the  $SL(2)$ -type is defined in terms of Tadic's classification of the unitary dual.

The reciprocity map for  $GL_n(F)$  is a bijection from the set of isomorphism classes of smooth irreducible representations of  $GL_n(F)$  to the set of isomorphism classes of  $n$ -dimensional Weil-Deligne representations (cf. [HT01] and [Hen00]). Applying the reciprocity map we observe that there is a natural way to extend the definition of the  $SL(2)$ -type to all smooth and irreducible representations of  $GL_n(F)$  (see Theorem 4.1 and Remark 1). The reciprocity map also allows the definition of base change with respect to any finite extension  $E$  of  $F$ . It is a map  $\text{bc}_{E/F}$  from isomorphism classes of smooth irreducible representation of  $GL_n(F)$  to isomorphism classes of smooth irreducible representation of  $GL_n(E)$  that is the 'mirror image' of restriction with

---

*Date:* December 7, 2008.

In this research the first named author is supported by THE ISRAEL SCIENCE FOUNDATION (grant No. 88/08).

respect to  $E/F$  of Weil-Deligne representations. The content of Theorem 6.1, our main result, is that for any smooth, irreducible and unitarizable representation  $\pi$  of  $GL_n(F)$  the representations  $\pi$  and  $\text{bc}(\pi)$  have the same  $SL(2)$ -type.

In [OS07b], [OS07a], [OS08] we studied the Klyachko models of smooth irreducible representations of  $GL_n(F)$ , that is, distinction of a representation with respect to certain subgroups that are a semi direct product of a unipotent and a symplectic group. Our results are also described in terms of Tadic's classification and depend, in fact, only on the  $SL(2)$ -type of a representation. For example, a smooth, irreducible and unitarizable representation  $\pi$  of  $GL_{2n}(F)$  is  $Sp_{2n}(F)$ -distinguished, i.e. it satisfies  $\text{Hom}_{Sp_{2n}(F)}(\pi, \mathbb{C}) \neq 0$ , if and only if the  $SL(2)$ -type of  $\pi$  consists entirely of even parts (and in this case  $\text{Hom}_{Sp_{2n}(F)}(\pi, \mathbb{C})$  is one dimensional [HR90, Theorem 2.4.2]). For unitarizable representations, our results on Klyachko models are reinterpreted here in terms of the  $SL(2)$ -type. As a consequence we show that Klyachko models are preserved under base-change with respect to any finite extension. In particular, we have

**Theorem 1.1.** *Let  $E/F$  be a finite extension of  $p$ -adic fields. A smooth, irreducible and unitarizable representation  $\pi$  of  $GL_{2n}(F)$  is  $Sp_{2n}(F)$ -distinguished if and only if  $\text{bc}_{E/F}(\pi)$  is  $Sp_{2n}(E)$ -distinguished.*

The rest of this note is organized as follows. After setting some general notation in Section 2, in Section 3 we recall the definition of the reciprocity map. In Section 4 we recall the definition of Venkatesh for the  $SL(2)$ -type of a unitarizable representation and extend it to all smooth irreducible representations. We recall (and reformulate in terms of the  $SL(2)$ -type) our results on symplectic (and more generally on Klyachko) models in Section 5. Our main observation Theorem 6.1 and its application to Klyachko models Corollary 6.1 are stated in Section 6 and proved in Section 7. The main theorem says that base change respects  $SL(2)$ -types and its corollary says that base change respects Klyachko types. Theorem 1.1 is a special case where the Klyachko type is purely symplectic.

## 2. NOTATION

Let  $F$  be a finite extension of  $\mathbb{Q}_p$  for some prime number  $p$  and let  $|\cdot|_F : F^\times \rightarrow \mathbb{C}^\times$  denote the standard absolute value normalized so that the inverse of uniformizers are mapped to the size of the residual field. Denote by  $W_F$  the Weil group of  $F$  and by  $I_F$  the inertia subgroup of  $W_F$ . We normalize the reciprocity map  $T_F : W_F \rightarrow F^\times$ , given by local class field theory, so that geometric Frobenius elements are mapped to

uniformizers. The map  $T_F$  defines an isomorphism from the abelianization  $W_F^{ab}$  of  $W_F$  to  $F^\times$  (this is the inverse of the Artin map). Let  $|\cdot|_{W_F} = |\cdot|_F \circ T_F$  denote the associated absolute value on  $W_F$ .

Denote by  $\mathbf{1}_\Omega$  the characteristic function of a set  $\Omega$ . Let  $\text{MS}_{\text{fin}}(\Omega)$  be the set of finite multisets of elements in  $\Omega$ , that is, the set of functions  $f : \Omega \rightarrow \mathbb{Z}_{\geq 0}$  of finite support. When convenient we will also denote  $f$  by  $\{\omega_1, \dots, \omega_1, \omega_2, \dots, \omega_2, \dots\}$  where  $\omega \in \Omega$  is repeated  $f(\omega)$  times. Let  $\mathcal{P} = \text{MS}_{\text{fin}}(\mathbb{Z}_{>0})$  be the set of partitions of positive integers and let

$$\mathcal{P}(n) = \{f \in \mathcal{P} : \sum_{k=1}^{\infty} k f(k) = n\}$$

denote the subset of partitions of  $n$ . For  $n, m \in \mathbb{Z}_{>0}$  let  $(n)_m = m \mathbf{1}_n = \{n, \dots, n\}$  be the partition of  $nm$  with ‘ $m$  parts of size  $n$ ’. Let  $\text{odd} : \mathcal{P} \rightarrow \mathbb{Z}_{\geq 0}$  be defined by

$$\text{odd}(f) = \sum_{k=0}^{\infty} f(2k+1),$$

i.e.  $\text{odd}(f)$  is the number of odd parts of the partition  $f$ .

### 3. RECIPROCITY AND BASE-CHANGE FOR $GL_n(F)$

**3.1. Weil-Deligne representations.** An  $n$ -dimensional *Weil-Deligne* representation is a pair  $((\rho, V), N)$  where  $(\rho, V)$  is an  $n$ -dimensional representation of  $W_F$  that decomposes as a direct sum of irreducible representations and  $N : V \rightarrow V$  is a linear operator such that

$$|w|_{W_F} N \circ \rho(w) = \rho(w) \circ N, \quad w \in W_F.$$

The map  $((\rho, V), N) \mapsto ([\rho], f)$ , where  $[\rho]$  denotes the isomorphism class of the  $n$ -dimensional representation  $(\rho, V)$  of  $W_F$  and  $f \in \mathcal{P}(n)$  is the partition of  $n$  associated to the Jordan decomposition of  $N$ , defines an injective map on isomorphism classes of Weil-Deligne representations. Denote its image by  $\mathcal{G}_F(n)$ . In this way we identify the set  $\mathcal{G}_F(n)$  with the set of isomorphism classes of  $n$ -dimensional Weil-Deligne representations. Let  $P_{F,n} : \mathcal{G}_F(n) \rightarrow \mathcal{P}(n)$  be the projection to the second coordinate. Let  $\mathcal{G}_F = \cup_{n=1}^{\infty} \mathcal{G}_F(n)$  be the set of isomorphism classes of all finite dimensional Weil-Deligne representations and let  $P_F : \mathcal{G}_F \rightarrow \mathcal{P}$  be the map such that  $P_F|_{\mathcal{G}_F(n)} = P_{F,n}$ .

**3.2. The local Langlands correspondence.** Let  $\mathcal{A}_F(n)$  be the set of isomorphism classes of smooth and irreducible representations of  $GL_n(F)$  and set  $\mathcal{A}_F = \cup_{n=1}^{\infty} \mathcal{A}_F(n)$ . For every  $\pi \in \mathcal{A}_F$  we denote by  $\omega_\pi$  the central character of (any representation in the isomorphism class of)  $\pi$ . Fix a non trivial additive character  $\psi$  of  $F$ . Due to Harris-Taylor

[HT01] and independently to Henniart [Hen00] there exists a unique sequence of bijections

$$\text{rec}_{F,n} : \mathcal{A}_F(n) \rightarrow \mathcal{G}_F(n)$$

for all  $n \geq 1$  satisfying the following properties:

$$(3.1) \quad \text{rec}_F(\chi) = \chi \circ T_F;$$

$$(3.2) \quad L(\pi_1 \times \pi_2, s) = L(\text{rec}_F(\pi_1) \otimes \text{rec}_F(\pi_2), s);$$

$$(3.3) \quad \epsilon(\pi_1 \times \pi_2, s, \psi) = \epsilon(\text{rec}_F(\pi_1) \otimes \text{rec}_F(\pi_2), s, \psi);$$

$$(3.4) \quad \det \circ \text{rec}_F(\pi) = \text{rec}_F(\omega_\pi);$$

$$(3.5) \quad \text{rec}_F(\pi^\vee) = \text{rec}_F(\pi)^\vee.$$

Here  $\chi \in \mathcal{A}_F(1)$ ,  $\pi, \pi_1, \pi_2 \in \mathcal{A}_F$ ,  $\pi^\vee$  is the contragredient of  $\pi$ ,  $\text{rec}_F(\pi)^\vee$  is the dual of  $\text{rec}_F(\pi)$  and  $\text{rec}_F : \mathcal{A}_F \rightarrow \mathcal{G}_F$  is such that  $\text{rec}_{F|\mathcal{A}_F(n)} = \text{rec}_{F,n}$ .

**3.3. Expressing  $\text{rec}_F$  in terms of  $\text{rec}_F^\circ$ .** Let  $\mathcal{A}_F^\circ(n) \subseteq \mathcal{A}_F(n)$  be the subset of isomorphism classes of supercuspidal representations and let  $\mathcal{G}_F^\circ(n) \subseteq \mathcal{G}_F(n)$  be the subset of isomorphism classes  $([\rho], f)$  such that  $\rho$  is irreducible and  $f = \mathbf{1}_n = \{n\}$ . The set  $\mathcal{G}_F^\circ(n)$  is identified with the set of isomorphism classes of irreducible and  $n$ -dimensional representations of  $W_F$ . It follows from the work of Harris-Taylor and independently of Henniart that there exists a unique sequence of bijections

$$\text{rec}_{F,n|\mathcal{A}_F^\circ(n)} = \text{rec}_{F,n}^\circ : \mathcal{A}_F(n) \rightarrow \mathcal{G}_F(n)$$

satisfying (3.1), (3.2), (3.3), (3.4) and (3.5). The work of Zelevinsky [Zel80] allows the extension of  $\text{rec}_F^\circ$  to the map  $\text{rec}_F$  on  $\mathcal{A}_F$ . This is also explained in [Hen85] and we now recall the construction of  $\text{rec}_F$  in terms of  $\text{rec}_F^\circ$ .

For  $s \in \mathbb{C}$  and every isomorphism class  $\varpi = [\pi] \in \mathcal{A}_F$  (resp.  $\varrho = ([\rho], f) \in \mathcal{G}_F$ ) let  $\varpi[s] = [\pi \otimes |\det|_F^s]$  (resp.  $\varrho[s] = ([\rho \otimes \cdot|_{W_F}^s], f)$ ). A *segment* in  $\mathcal{A}_F$  (resp.  $\mathcal{G}_F^\circ$ ) is a set of the form

$$\Delta[\sigma, r] = \left\{ \sigma\left[\frac{1-r}{2}\right], \sigma\left[\frac{3-r}{2}\right], \dots, \sigma\left[\frac{r-1}{2}\right] \right\}$$

(resp.

$$\Delta[\rho, r] = \left\{ \rho\left[\frac{1-r}{2}\right], \rho\left[\frac{3-r}{2}\right], \dots, \rho\left[\frac{r-1}{2}\right] \right\})$$

for some  $\sigma \in \mathcal{A}_F$  (resp.  $\rho \in \mathcal{G}_F^\circ$ ) and  $r \in \mathbb{Z}_{>0}$ . Let  $\mathcal{S}$  (resp.  $\mathcal{S}'$ ) denote the set of all segments in  $\mathcal{A}_F$  (resp.  $\mathcal{G}_F^\circ$ ) and let  $\mathcal{O} = \text{MS}_{\text{fin}}(\mathcal{S})$  (resp.  $\mathcal{O}' = \text{MS}_{\text{fin}}(\mathcal{S}')$ ). The bijection  $\text{rec}_F^\circ : \mathcal{A}_F \rightarrow \mathcal{G}_F^\circ$  defines a bijection  $\text{rec}_F^\circ : \mathcal{S} \rightarrow \mathcal{S}'$  given by  $\text{rec}_F^\circ(\Delta[\sigma, r]) = \Delta[\text{rec}_F^\circ(\sigma), r]$  and a bijection  $\text{rec}_F^\circ : \mathcal{O} \rightarrow \mathcal{O}'$  given by  $\text{rec}_F^\circ(a)(\text{rec}_F^\circ(\Delta)) = a(\Delta)$ ,  $\Delta \in \mathcal{S}$ .

In [Zel80, Section 6.5] Zelevinsky defines a bijection  $a \mapsto \langle a \rangle$  from  $\mathcal{O}$  to  $\mathcal{A}_F$ . The Zelevinsky involution is defined in [Zel80, Section 9.12] as an involution on the Grothendick group associated with  $\mathcal{A}_F$ . It is proved by Aubert [Aub95], [Aub96] and independently by Procter [Pro98] that the Zelevinsky involution restricts to a bijection from  $\mathcal{A}_F$  to itself that we denote by  $\pi \mapsto \pi^t$ . In [Zel80, Section 10.2] Zelevinsky defines a bijection  $\tau : \mathcal{O}' \rightarrow \mathcal{G}_F$  as follows. For a segment  $\Delta[\rho, r] \in \mathcal{S}'$  where  $\rho \in \mathcal{G}_F^\circ(t)$  let

$$\tau(\Delta[\rho, r]) = (\oplus_{i=1}^r \rho, (r)_t)$$

and for  $a' \in \mathcal{O}'$  set

$$\tau(a') = \oplus_{\Delta' \in \mathcal{O}'} \tau(\Delta')$$

where for  $([\rho_1], f_1), \dots, ([\rho_m], f_m) \in \mathcal{G}_F$  the direct sum is given by

$$([\rho_1], f_1) \oplus \dots \oplus ([\rho_m], f_m) = ([\rho_1 \oplus \dots \oplus \rho_m], f_1 + \dots + f_m).$$

The reciprocity map  $\text{rec}_F$  is given by

$$\text{rec}_F(\langle a \rangle^t) = \tau(\text{rec}_F^\circ(a)), \quad a \in \mathcal{O}.$$

#### 4. THE $SL(2)$ -TYPE OF A REPRESENTATION

Denote by  $\mathcal{A}_F^u(n)$  the subset of  $\mathcal{A}_F(n)$  consisting of all isomorphism classes of unitarizable representations and let  $\mathcal{A}_F^u = \cup_{n=1}^\infty \mathcal{A}_F^u(n)$ . For  $[\pi_1], \dots, [\pi_m] \in \mathcal{A}_F$  we denote by  $\pi_1 \times \dots \times \pi_m$  the representation parabolically induced from  $\pi_1 \otimes \dots \otimes \pi_m$  and by  $[\pi_1] \times \dots \times [\pi_m]$  its isomorphism class.

For  $\sigma \in \mathcal{A}_F^\circ$  and integers  $n, r > 0$  let

$$\delta[\sigma, n] = \langle \Delta[\sigma, n] \rangle^t,$$

$$a(\sigma, n, r) = \{ \Delta[\sigma[\frac{1-r}{2}], n], \Delta[\sigma[\frac{3-r}{2}], n], \dots, \Delta[\sigma, n](\frac{r-1}{2}) \} \in \mathcal{O}$$

and

$$U(\delta[\sigma, n], r) = \langle a(\sigma, n, r) \rangle.$$

Tadic's classification of the unitary dual of  $GL_n(F)$  [Tad86] implies that if  $\sigma \in \mathcal{A}_F^\circ \cap \mathcal{A}_F^u$  then  $U(\delta[\sigma, n], r) \in \mathcal{A}_F^u$  and that for any  $\pi \in \mathcal{A}_F^u$  there exist  $\sigma_1, \dots, \sigma_m \in \mathcal{A}_F^\circ$  and integers  $n_1, \dots, n_m, r_1, \dots, r_m > 0$  such that

$$(4.1) \quad \pi = U(\delta[\sigma_1, n_1], r_1) \times \dots \times U(\delta[\sigma_m, n_m], r_m).$$

It further follows from [Tad95, Lemma 3.3] that

$$(4.2) \quad U(\delta[\sigma, n], r)^t = U(\delta[\sigma, r], n).$$

The  $SL(2)$  of a representation  $\pi \in \mathcal{A}_F^u$  of the form (4.1) is defined in [Ven05, Definition 1] to be the partition

$$(4.3) \quad \{(r_1)_{n_1}, \dots, (r_m)_{n_m}\}.$$

**Theorem 4.1.** *The  $SL(2)$ -type of a representation  $\pi \in \mathcal{A}_F^u$  equals  $P_F(\text{rec}_F(\pi^t))$ .*

*Remark 1.* Theorem 4.1 allows us to define the  $SL(2)$ -type of any  $\pi \in \mathcal{A}_F$  by the formula  $P_F(\text{rec}_F(\pi^t))$ . Note further that given a reciprocity map (local Langlands conjecture), this provides a recipe to define the  $SL(2)$ -type of an irreducible representation for any reductive group!

*Proof.* Based on Tadic's classification of the unitary dual of  $GL_n(F)$ , the proof of Theorem 4.1 is merely a matter of following the definitions. For convenience, we provide the proof. The key is in the following simple observations.

**Lemma 4.1.** *Let  $\pi \in \mathcal{A}_F^u$  be of the form (4.1). Then*

$$(4.4) \quad \text{rec}_F(\pi) = \bigoplus_{i=1}^m \bigoplus_{j=1}^{r_i} \tau(\Delta[\sigma_i[\frac{r_i+1}{2} - j], n_i])$$

and

$$(4.5) \quad \pi^t = U(\delta[\sigma_1, r_1], n_1) \times \dots \times U(\delta[\sigma_m, r_m], n_m) \in \mathcal{A}_F^u.$$

*Proof.* Let  $a_i = a(\sigma_i, r_i, n_i)$ . It follows from (4.2) that

$$\pi = \langle a_1 \rangle^t \times \dots \times \langle a_m \rangle^t = (\langle a_1 \rangle \times \dots \times \langle a_m \rangle)^t$$

and since  $t$  is an involution on  $\mathcal{A}_F$  that  $\langle a_1 \rangle \times \dots \times \langle a_m \rangle \in \mathcal{A}_F$ . Thus, it follows from [Zel80, Proposition 8.4] that  $\langle a_1 \rangle \times \dots \times \langle a_m \rangle = \langle a_1 + \dots + a_m \rangle$ . In other words  $\pi = \langle a_1 + \dots + a_m \rangle^t$  and therefore by definition

$$\text{rec}_F(\pi) = \tau(\text{rec}_F^\circ(a_1 + \dots + a_m)) = \bigoplus_{i=1}^m \tau(\text{rec}_F^\circ(a_i)).$$

The identity (4.4) now follows from the definition of  $\tau(\text{rec}_F^\circ(a_i))$ . Note that (4.2) implies that

$$\pi^t = U(\delta[\sigma_1, r_1], n_1) \times \dots \times U(\delta[\sigma_m, r_m], n_m)$$

and the classification of Tadic therefore implies that  $\pi^t \in \mathcal{A}_F^u$ . Thus we get (4.5).  $\square$

Applying (4.4) to  $\pi^t$  and comparing with (4.3) Theorem 4.1 follows from the definitions.  $\square$

From now on for every  $\pi \in \mathcal{A}_F$  we denote by

$$(4.6) \quad \mathcal{V}(\pi) = P_F(\text{rec}_F(\pi^t))$$

the  $SL(2)$ -type of  $\pi$ .

## 5. KLYACHKO MODELS

For positive integers  $r$  and  $k$  denote by  $U_r$  the subgroup of upper triangular unipotent matrices in  $GL_r(F)$  and by  $Sp_{2k}(F)$  the symplectic group in  $GL_{2k}(F)$ . Fix a decomposition  $n = r + 2k$ . Let

$$H_{r,2k} = \left\{ \begin{pmatrix} u & X \\ 0 & h \end{pmatrix} : u \in U_r, X \in M_{r \times 2k}(F), h \in Sp_{2k}(F) \right\}.$$

Let  $\psi$  be a non trivial character of  $F$ . For  $u = (u_{i,j}) \in U_r$  let

$$\psi_r(u) = \psi(u_{1,2} + \cdots + u_{r-1,r})$$

and let  $\psi_{r,2k}$  be the character of  $H_{r,2k}$  defined by

$$\psi_{r,2k} \left( \begin{pmatrix} u & X \\ 0 & h \end{pmatrix} \right) = \psi_r(u).$$

We refer to the space

$$\mathcal{M}_{r,2k} = \text{Ind}_{H_{r,2k}}^{GL_n(F)}(\psi_{r,2k})$$

as a *Klyachko model* for  $GL_n(F)$ . Here  $\text{Ind}$  denotes the functor of non-compact smooth induction.

In [OS08, Corollary 1] we showed that for any  $\pi \in \mathcal{A}_F^u(n)$  there exists a unique decomposition

$$n = r(\pi) + 2k(\pi)$$

such that

$$\text{Hom}_{GL_n(F)}(\pi, \mathcal{M}_{r(\pi), 2k(\pi)}) \neq 0$$

and that in fact  $\dim_{\mathbb{C}}(\text{Hom}_{GL_n(F)}(\pi, \mathcal{M}_{r(\pi), 2k(\pi)})) = 1$ .

**Definition 1.** For  $\pi \in \mathcal{A}_F^u$ , the *Klyachko type* of  $\pi$  is the ordered pair  $(r(\pi), 2k(\pi))$ .

In fact, for  $\mathcal{A}_F^u$  [OS07a, Theorem 8] provides a receipt in order to read the Klyachko type off Tadic's classification. Based on (4.3), our results can be reinterpreted by the formula

$$(5.1) \quad r(\pi) = \text{odd}(\mathcal{V}(\pi)), \quad \pi \in \mathcal{A}_F^u.$$

## 6. BASE CHANGE-THE MAIN RESULTS

Let  $E$  be a finite extension of  $F$ . Denote by  $\text{res}_{E/F,n} : \mathcal{G}_F(n) \rightarrow \mathcal{G}_E(n)$  the map defined by  $\text{res}_{E/F,n}([\rho], f) = ([\rho|_{W_E}], f)$ . For  $n \geq 1$  the *base change*  $\text{bc}_{E/F}(\pi) \in \mathcal{A}_E(n)$  of  $\pi \in \mathcal{A}_F(n)$  is defined by

$$\text{rec}_E(\text{bc}_{E/F}(\pi)) = \text{res}_{E/F}(\text{rec}_F(\pi)).$$

**Theorem 6.1.** *Let  $E/F$  be a finite extension of  $p$ -adic fields and let  $\pi$  be a smooth, irreducible and unitarizable representation of  $GL_n(F)$ . Then  $\mathrm{bc}_{E/F}(\pi)$  is a smooth, irreducible and unitarizable representation of  $GL_n(E)$  and*

$$\mathcal{V}(\pi) = \mathcal{V}(\mathrm{bc}_{E/F}(\pi)),$$

*i.e.  $\pi$  and  $\mathrm{bc}_{E/F}(\pi)$  have the same  $SL(2)$ -type.*

As a consequence we have the following.

**Corollary 6.1.** *Under the assumptions of Theorem 6.1 we have*

$$r(\pi) = r(\mathrm{bc}_{E/F}(\pi)),$$

*i.e.  $\pi$  and  $\mathrm{bc}_{E/F}(\pi)$  have the same Klyachko type.*

Corollary 6.1 is straightforward from Theorem 6.1 and (5.1).

## 7. PROOF OF THE MAIN RESULT

**Lemma 7.1.** *Let  $E/F$  be a finite extension. For  $\sigma \in \mathcal{A}_F^\circ \cap \mathcal{A}_F^u$  there exist  $\sigma_1, \dots, \sigma_m \in \mathcal{A}_E^\circ \cap \mathcal{A}_E^u$  such that*

$$\mathrm{bc}_{E/F}(\sigma) = \sigma_1 \times \cdots \times \sigma_m.$$

*Proof.* Recall that a representation in  $\mathcal{A}_F^\circ$  is unitarizable if and only if its central character is unitary. Let  $\rho$  be the irreducible representation of  $W_F$  such that  $\mathrm{rec}_F(\sigma) = ([\rho], \mathbf{1}_n)$ . It follows from (3.4) that  $\rho$  has a unitary central character and therefore it has a unitary structure. Thus, the restriction  $\rho|_{W_E}$  to  $W_E$  also has a unitary structure and therefore each of its irreducible components has a unitary central character. The lemma follows by applying (4.4) to  $\mathrm{res}_{E/F}(\mathrm{rec}_F(\sigma))$ .  $\square$

**Proposition 7.1.** *Let  $E/F$  be a finite extension and let  $\pi \in \mathcal{A}_F^u$  then  $\mathrm{bc}(\pi) \in \mathcal{A}_E^u$  and*

$$(7.1) \quad \mathrm{bc}_{E/F}(\pi^t) = \mathrm{bc}_{E/F}(\pi)^t.$$

*Proof.* Let  $\pi \in \mathcal{A}_F^u$  be of the form (4.1). By Lemma 7.1 there exist  $\sigma_{i,k} \in \mathcal{A}_E^\circ$ ,  $i = 1, \dots, m$ ,  $k = 1, \dots, t_i$  such that

$$\mathrm{bc}_{E/F}(\sigma_i) = \sigma_{i,1} \times \cdots \times \sigma_{i,t_i}.$$

Let  $\rho_i = \mathrm{rec}_F^\circ(\sigma_i)$  and  $\rho_{i,k} = \mathrm{rec}_E^\circ(\sigma_{i,k})$ . Thus,

$$\mathrm{res}_{E/F}(\rho_i) = \bigoplus_{k=1}^{t_i} \rho_{i,k}.$$

It follows from (4.4) that

$$(7.2) \quad \mathrm{res}_{E/F}(\mathrm{rec}_F(\pi)) = \bigoplus_{i=1}^m \bigoplus_{j=1}^{r_i} \bigoplus_{k=1}^{t_i} \tau(\Delta[\sigma_{i,k}[\frac{r_i+1}{2} - j], n_i]).$$



On the other hand, let

$$\Pi = \times_{i=1}^m \times_{k=1}^{t_i} U(\delta[\sigma_{i,k}, n_i], r_i)$$

Since  $\pi \in \mathcal{A}_F^u$ , the classification of Tadic implies that  $\Pi \in \mathcal{A}_E^u$  and by (4.4) applied to  $E$  instead of  $F$  we have

$$(7.3) \quad \text{rec}_E(\Pi) = \oplus_{i=1}^m \oplus_{j=1}^{r_i} \oplus_{k=1}^{t_i} \tau(\Delta[\sigma_{i,k}[\frac{r_i+1}{2} - j], n_i]).$$

Comparing (7.2) with (7.3) we obtain that  $\Pi = \text{bc}_{E/F}(\pi)$  and in particular that  $\text{bc}_{E/F}(\pi) \in \mathcal{A}_E^u$ . Applying this to  $\pi^t$  expressed by (4.5) gives

$$\text{bc}_{E/F}(\pi^t) = \times_{i=1}^m \times_{k=1}^{t_i} U(\delta[\sigma_{i,k}, r_i], n_i).$$

Applying (4.5) now to  $\text{bc}_{E/F}(\pi)^t$  we obtain the identity (7.1).  $\square$

It is straightforward from the definitions that

$$(7.4) \quad P_F(\text{rec}_F(\pi)) = P_E(\text{rec}_E(\text{bc}_{E/F}(\pi))), \quad \pi \in \mathcal{A}_F.$$

For  $\pi \in \mathcal{A}_F^u$ , applying (7.4) to  $\pi^t$  and then (7.1) we get that

$$P_F(\text{rec}_F(\pi^t)) = P_E(\text{rec}_E(\text{bc}_{E/F}(\pi)^t)).$$

The identity  $\mathcal{V}(\pi) = \mathcal{V}(\text{bc}_{E/F}(\pi))$  is now immediate from (4.6). This completes the proof of Theorem 6.1.

## REFERENCES

- [Aub95] Anne-Marie Aubert. Dualité dans le groupe de Grothendieck de la catégorie des représentations lisses de longueur finie d'un groupe réductif  $p$ -adique. *Trans. Amer. Math. Soc.*, 347(6):2179–2189, 1995.
- [Aub96] Anne-Marie Aubert. Erratum: “Duality in the Grothendieck group of the category of finite-length smooth representations of a  $p$ -adic reductive group” [Trans. Amer. Math. Soc. **347** (1995), no. 6, 2179–2189; [Aub95]]. *Trans. Amer. Math. Soc.*, 348(11):4687–4690, 1996.
- [Hen85] Guy Henniart. Le point sur la conjecture de Langlands pour  $GL(N)$  sur un corps local. In *Séminaire de théorie des nombres, Paris 1983–84*, volume 59 of *Progr. Math.*, pages 115–131. Birkhäuser Boston, Boston, MA, 1985.
- [Hen00] Guy Henniart. Une preuve simple des conjectures de Langlands pour  $GL(n)$  sur un corps  $p$ -adique. *Invent. Math.*, 139(2):439–455, 2000.
- [HR90] Michael J. Heumos and Stephen Rallis. Symplectic-Whittaker models for  $GL_n$ . *Pacific J. Math.*, 146(2):247–279, 1990.
- [HT01] Michael Harris and Richard Taylor. *The geometry and cohomology of some simple Shimura varieties*, volume 151 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001. With an appendix by Vladimir G. Berkovich.
- [OS07a] Omer Offen and Eitan Sayag. Global mixed periods and local klyachko models for the general linear group. *Internation Math. Reaseach Notices*, 2007:article ID rnm136, 25 pages, doi:10.1093/imrn/rnm136, 2007.

- [OS07b] Omer Offen and Eitan Sayag. On unitary representations of  $GL_{2n}$  distinguished by the symplectic group. *J. Number Theory*, 125(2):344–355, 2007.
- [OS08] Omer Offen and Eitan Sayag. Uniqueness and disjointness of klyachko models. *J. of Functional Analysis*, 254:2846–2865, 2008.
- [Pro98] Kerrigan Procter. Parabolic induction via Hecke algebras and the Zelevinsky duality conjecture. *Proc. London Math. Soc. (3)*, 77(1):79–116, 1998.
- [Tad86] Marko Tadić. Classification of unitary representations in irreducible representations of general linear group (non-Archimedean case). *Ann. Sci. École Norm. Sup. (4)*, 19(3):335–382, 1986.
- [Tad95] M. Tadić. On characters of irreducible unitary representations of general linear groups. *Abh. Math. Sem. Univ. Hamburg*, 65:341–363, 1995.
- [Ven05] Akshay Venkatesh. The Burger-Sarnak method and operations on the unitary dual of  $GL(n)$ . *Represent. Theory*, 9:268–286 (electronic), 2005.
- [Zel80] A. V. Zelevinsky. Induced representations of reductive  $\mathfrak{p}$ -adic groups. II. On irreducible representations of  $GL(n)$ . *Ann. Sci. École Norm. Sup. (4)*, 13(2):165–210, 1980.